






Catalogue of tree microhabitats

Reference field list



This catalogue is available for download:
integrateplus.org

Recommended citation: Kraus, D., Bütler, R., Krumm, F., Lachat, T., Larrieu, L., Mergner, U., Paillet, Y., Rydkvist, T., Schuck, A., and Winter, S. 2016. Catalogue of tree microhabitats – Reference field list. Integrate+ Technical Paper. 16p.

Illustrations: Lisa Apfelbacher

Photos: Daniel Kraus

Disclaimer: The present catalogue is one of the outputs of the demonstration project '*Establishing a European network of demonstration sites for the integration of biodiversity conservation into forest management*' supported by the German Federal Ministry of Food and Agriculture (BMEL). The views expressed in this publication are those of the authors and do not necessarily represent those of the European Forest Institute.

European Forest Institute, 2016

Large quantities of deadwood and a high density of old microhabitat-bearing trees are characteristic elements of natural forests, especially of the old-growth phases. These are often absent or rare in managed forests, even in forests under close-to-nature management. Yet, an important share of forest biodiversity is strictly or primarily dependent on such elements for their survival, especially 'saproxylic' species, those are species depending on deadwood.



Tree related microhabitats are therefore recognised as important substrates and structures for biodiversity in forests. The retention of both existing and future tree microhabitats is thus one important aspect to take in to consideration in forest management. Giving tree microhabitats increased attention will help sustain and increase the habitat value for biodiversity also in managed forests .

This reference field list is developed to support training exercises conducted in Integrate+ Marteloscope sites. It aims at supporting forest managers, inventory personnel and other groups in identifying and describing tree microhabitats in the course of such exercises. It can also find use as illustrative material in forest education and as background documentation for other training events and field excursions.



Saproxyllic microhabitats	Description	Type	Code	Illustrations
---------------------------	-------------	------	------	---------------

Cavities

At least three in the trunk connected woodpecker breeding cavities. If this cannot be checked: three cavity openings within two meters.	Woodpecker "flute" / cavity string	CV15	
Trunk and mould cavities			
Trunk cavity with mould, cavity bottom has ground contact thus soil humidity enters the cavity hole. Note that the cavity entrance can be higher at the trunk.	ø ≥ 10 cm (ground contact)	CV21	
	ø ≥ 30 cm (ground contact)	CV22	
Mould containing trunk cavity without ground contact.	ø ≥ 10 cm	CV23	
	ø ≥ 30 cm	CV24	
			
Semi-open trunk cavity with or without mould, cavity chamber is not completely protected from surrounding microclimate and precipitation may enter the interior. Note that the cavity entrance can be higher at the trunk.	ø ≥ 30 cm / semi-open	CV25	
Large trunk cavity with open top and with or without ground contact.	ø ≥ 30 cm /open top	CV26	

Illustrations	Code	Type	Description	Saproxyllic microhabitats
---------------	------	------	-------------	---------------------------

CV3

Brach holes				
	CV31	$\varnothing \geq 5\text{ cm}$	Rot-holes originating from branch breakage at trunk when fungal decay of wood is progressing faster than occlusion of wound.	
	CV32	$\varnothing \geq 10\text{ cm}$		
				
	CV33	Hollow branch, $\varnothing \geq 10\text{ cm}$	Hollow more or less horizontal branch developing from breakage. Provides tubular shelter from surrounding microclimate.	

CV4

Dendrotelms and water-filled holes			
	CV41	$\varnothing \geq 3\text{ cm}$ / trunk base	The entrance diameter is the same as the chamber's interior. Cup-shaped concavities that, due to their form, retain precipitation for a certain period of time (until it dries out).
	CV42	$\varnothing \geq 15\text{ cm}$ / trunk base	
	CV43	$\varnothing \geq 5\text{cm}$ / crown	The entrance diameter is the same as the chamber's interior. Cup-shaped concavities that, due to their form, retain precipitation for a certain period of time (until it dries out).
	CV44	$\varnothing \geq 15\text{ cm}$ / crown	
			

Saproxyllic microhabitats	Description	Type	Code	Illustrations
---------------------------	-------------	------	------	---------------

Cavities

Insect galleries and bore holes				
The entrance or exit diameter is the same as the interior's hole diameter. A bore hole net of xylophagous insects indicates a wood hole system. An insect gallery is a complex system of holes and chambers created by one or more insect species within the trunk.	Gallery with single small bore holes		CV51	
	Large bore hole ø ≥ 2 cm		CV52	


Injuries and wounds



Bark loss / exposed sapwood				
Loss of trunk bark thus sapwood is exposed; caused e.g. by felling, natural falling of trees, rock fall. At the trunk base, bark loss may also be caused by skidding of logs, rodents, woodpecker sloughing.	Bark loss 25- 600 cm ² , decay stage < 3		IN 11	
	Bark loss > 600 cm ² , decay stage < 3		IN12	
	Bark loss 25- 600 cm ² , Decay stage = 3		IN13	
	Bark loss > 600 cm ² , decay stage = 3		IN14	
Exposed heartwood / trunk and crown breakage				
The tree has broken off at the trunk level, in a living tree. The tree is still alive and is developing a secondary crown with parts of the trunk decaying near the injury: the tree combines large decaying wood with xylem and phloem flux.	Broken trunk, ø ≥ 20 cm at the broken end		IN21	

CV5

IN1

IN2

Illustrations	Code	Type	Description	Saprophytic microhabitats
 	IN22	Broken tree crown / fork Exposed wood ≥ 300 cm ²	Exposed heartwood through the fork insertion breakage into the trunk - the rot initiates decaying substrate on the living tree.	Injuries and wounds
 	IN23	Broken limb, $\varnothing \geq 20$ cm at the broken end	A 1 st order branch has broken off. The tree is still alive. The injury provides a large entry gate for organisms and may develop into a cavity (rot hole) with xylem and phloem flux.	
	IN24	Splintered stem, $\varnothing \geq 20$ cm at the broken end	At wind breakage, trunk has splintered with several long splinters due to high force: splintered wound provides specific ecological conditions.	
Cracks and scars				
	IN31	Length ≥ 30 cm; width > 1 cm; depth > 10 cm	Line-shaped injury (cleft) through the bark into the sapwood, exposing cambium and sapwood (not to be recorded if injury has occluded).	
	IN32	Length ≥ 100 cm; width > 1 cm; depth > 10 cm		

Saproxyltic microhabitats	Description	Type	Code	Illustrations
Injuries and wounds	Bark loss and crack caused by lightning strike exposing the sapwood (not recorded when new bark has closed the scar).	Lightning scar	IN33	
	Fire scars at the lower trunk usually have a triangular shape and are located at the base of the tree on the leeward trunk side. Fire scars are associated with charred wood and eventually resin flow on exposed sapwood or bark.	Fire scar, $\geq 600 \text{ cm}^2$	IN34	
Bark		Bark		
	Space between bark and sapwood forming a shelter (open at the bottom).	Bark shelter, width > 1 cm; depth > 10 cm; height > 10 cm	BA11	
	Space between bark and sapwood forming a pocket (open at the top), eventually containing mould.	Bark pocket, width > 1 cm; depth > 10 cm; height > 10 cm	BA12	
	Coarse and fissured bark, sometimes tree species specific.	Coarse bark	BA21	

Illustrations	Code	Type	Description	Saproxyllic microhabitats
---------------	------	------	-------------	---------------------------

DE1



Dead branches and limbs / crown deadwood

DE11	ø 10 - 20 cm, ≥ 50 cm, sun exposed	Smaller sized (> 10 cm diameter) decaying wood, often horizontal or at a skewed angle, often in the shadow of the remaining canopy; in contact with living wood (xylem and phloem flow).
DE12	ø > 20 cm, ≥ 50 cm, sun exposed	
DE13	ø 10 - 20 cm, ≥ 50 cm, not sun exposed	
DE14	ø > 20 cm, ≥ 50 cm, not sun exposed	
DE15	Dead top ø ≥ 10 cm	

Deadwood

Illustrations	Code	Type	Description	Epixylic microhabitats
---------------	------	------	-------------	------------------------

GR1



Root buttress cavities

GR11	ø ≥ 5 cm	Natural cavity at the base of the tree trunk formed by the tree roots. May be densely covered with bryophytes. No wound or rot-hole.
GR12	ø ≥ 10 cm	
GR13	Trunk cleavage length ≥ 30 cm	Cleft formed by tree growth, no wound or open crack. Enclosure located higher at the trunk and therefore not part of the root buttress.

Deformation / growth form

Epixylic microhabitats	Description	Type	Code	Illustrations
------------------------	-------------	------	------	---------------


Deformation / growth form

Witches broom

Dense agglomeration of twigs caused by a parasite (such the fungi *Melampsorella caryophylacerum* or *Taphrina betulina*) or hemiparasite (genus *Arceuthobium*, *Viscaceae*).

Witches broom,
ø > 50 cm

GR21



GR2

Dense agglomeration of shoots on the trunk or branches of a tree. They originate from latent buds visible on the tree or can be submerged under the bark as epicormic buds.

Water sprout

GR22



Cankers and burrs

Proliferation of cell growth with rough bark and bark damage at the canker surface.

Cancerous growth,
ø > 20 cm

GR31



GR3

Decayed canker exposing necrotic tissue, e.g. caused by *Nectria spec.* on beech.

Decayed canker,
ø > 20 cm

GR32



Illustrations	Code	Type	Description	Epixylic microhabitats
---------------	------	------	-------------	------------------------

Fruiting bodies fungi



EP11 Annual polypores, $\varnothing > 5\text{cm}$

Fruiting bodies of polypores at the tree trunk lasting some weeks. European polypores only have one layer of tubes and are mostly of tough elastic soft consistency (no woody parts). Many species do not develop fruiting bodies each year. Main annual genera are *Abortiporus*, *Amylocystis*, *Bjerkandera*, *Bondar-zewia*, *Cerrena*, *Climacocystis*, *Fistulina*, *Gloeophyllum*, *Grifola*, *Haploporus*, *Inonotus*, *Ischnoderma*, *Laetiporus*, *Leptoporus*, *Meripilus*, *Oligoporus*, *Oxyporus*, *Perenniporia* pp, *Phaeolus*, *Piptoporus*, *Podofomes*, *Polyporus*, *Pycnoporus*, *Spongipellis*, *Stereum*, *Trametes*, *Trichaptum*, *Tyromyces* (underlined genera known to host a wide diversity / rare invertebrates).



EP12 Perennial polypores, $\varnothing > 10\text{cm}$

Woody, or at least tough fruiting bodies, showing distinct annual layers in the tube layer. Perennial fruiting bodies of the fungi indicating trunk decay caused by white rot (e.g. *Fomes fomentarius* (L. ex Fr.) Fr.) and brown rot (e.g. *Fomitopsis pinicola* (Swartz ex Fr.) Karst.). Main perennial genera are *Fomitopsis* pp, *Fomes*, *Perreniporia* pp, *Oxyporus*, *Ganoderma* pp, *Phellinus*, *Daedalea*, *Haploporus*, *Heterobasidion*, *Hexagonia*, *Laricifomes*, *Daedaleopsis* (underlined genera known to host a wide diversity / rare invertebrates).



EP13 Pulpy agaric, $\varnothing > 5\text{cm}$

Large, thick and pulpy or rather fleshy fruiting body of gill-bearing fungus (order Agaricales) - an agaric is a type of fungal fruiting body characterized by the presence of a pileus (cap) that is clearly differentiated from the stipe (stalk), with lamellae (gills) on the under-side of the pileus. "Agaric" can also refer to a basidiomycete species characterized by an agaric-type fruiting body. Examples: *Armillaria*, *Pleurotus*, *Megacollybia*, large *Pluteus* bear many arthropods and also parasitic fungi. The fruiting body remains generally several weeks.

Epiphytes

Epixylic microhabitats	Description	Type	Code	Illustrations
Epiphytes	Fungi cover of large tough hemispheric dark fungus looking like a lump of coal. Genus examples are <i>Daldinia</i> and <i>Hypoxylon</i> .	Large ascomycetes, $\varnothing > 5 \text{ cm}$	EP14	
	Myxomycetes			
	Amoeboid slime mould which forms moving plasmodium looking like gelatinous mass when fresh.	Myxomycetes, $\varnothing > 5 \text{ cm}$	EP21	
	Epiphytic crypto- and phanerogams			
	Tree trunk covered by mosses and liverworts.	Epiphytic bryophytes coverage > 25 %	EP31	
	Tree trunk covered by foliose and fruticose lichens (lichens in combination with bryophytes).	Epiphytic foliose and fruticose lichens, coverage > 25 %	EP32	
	Lianas and other climbing plants cover the trunk surface (e.g. <i>Hedera helix</i> , <i>Clematis vitalba</i>).	Lianas, coverage > 25 %	EP33	

EP2

EP3

Illustrations	Code	Type	Description	Epixylic microhabitats
	EP34	Epiphytic ferns, > 5 fronds	Epiphytic ferns on trunk and large branches, often associated with bryophytes.	Epiphytes
	EP35	Mistletoe	Occurrence of these epiphytic and hemiparasitic plant species in the tree crown (<i>Viscum spp.</i> , <i>Arceuthobium spp.</i> , <i>Amyena spp.</i> , <i>Loranthus spp.</i>).	
Nests				
	NE11	Large vertebrate nest, ø > 80 cm	Structures built by big raptors (eagles, black or white stork, grey heron) to hold eggs, offspring, or occasionally the animal itself. They may be composed of organic material such as twigs, grass, and leaves, and are located on branches, forks or witch brooms.	Nests
	NE12	Small vertebrate nest, ø > 10 cm	Nests built by small bird species, dormouse, mouse or squirrel.	
	NE21	Invertebrate nest	Larval nest of the pine processionary moth <i>Thaumetopoea pityocampa</i> , nest of the wood ant <i>Lasius fuliginosus</i> and of feral bees in tree trunk.	
Sap and resin run				
	OT11	Sap flow, > 50 cm	Fresh significant flow of sap, mainly at deciduous tree species.	Other

NE1

OT1

Epixylic microhabitats	Description	Type	Code	Illustrations
Other	Fresh significant flow of resin, at coniferous tree species.	Resin flow and pockets, > 50 cm	OT12	
	Result of micro-pedogenesis from epiphytic mosses, lichens or algae and necrosed old bark, it also depends on debris and litter fall from the crowns and bark.	Microsoil		
		Crown microsoil	OT21	
		Bark microsoil	OT22	

OT2



Integrate+ is a demonstration project funded by the German BMEL to establish a European network of demonstration sites for the integration of biodiversity conservation into forest management.

The Integrate+ project runs from December 2013 to December 2016 and builds on a partner network from research and practice with a focus on implementation of integrative management and enhancing transnational exchange of experiences.



European Forest Institute
Regional Office EFICENT
Wonnhaldestr. 4
79100 Freiburg, Germany

www.integrateplus.org
info@integrateplus.org